

Biochemistry test report



Patient:	BAILEY	Species:	Canine	Patient ID:	260115001
Client:	SALVA	Gender:	Male	Sample No.:	04
Doctor:		Age:	2Y	Time of analysis:	2026/01/15 09:49

	Item	Current result	Ref. Ranges
Protein	TP	↓ 26.7	g/L 53.1-79.2
Protein	ALB	↓ 7.2	g/L 23.4-40.0
Protein	GLOB	↓ 19.4	g/L 25.4-52.0
Protein	A/G	0.4	
Liver and gallbladder	ALT	45.2	U/L 10.1-100.3
Liver and gallbladder	AST	↑ 61.8	U/L 0.0-51.7
Liver and gallbladder	AST/ALT	1.37	
Liver and gallbladder	ALP	↑ I- 493.1	U/L 15.5-212.0
Liver and gallbladder	GGT	↑ 25.7	U/L 0.0-15.9
Liver and gallbladder	TBIL	↑ 80.68	μmol/L 0.00-15.00
Liver and gallbladder	TBA	↑ 43.1	μmol/L 0.0-30.0
Pancreas	AMY	607.0	U/L 397.7-1285.1
Kidneys	BUN	↑ 18.59	mmol/L 2.50-9.77
Kidneys	CREA	111.80	μmol/L 20.00-123.70
Kidneys	BUN/CREA	41.1	
Cardiovasc./Muscle	CK	↑ 507.5	U/L 66.4-257.5
Cardiovasc./Muscle	LDH	77.4	U/L 0.0-143.6
Energy metabolism	GLU	4.56	mmol/L 3.80-7.50
Energy metabolism	TC	↑ 9.24	mmol/L 2.67-8.38
Energy metabolism	TG	↑ 2.93	mmol/L 0.10-1.30
Minerals	Ca	↓ 1.40	mmol/L 2.10-2.97
Minerals	PHOS	↑ 3.01	mmol/L 0.80-2.20
Minerals	CaxP	4.21	mmol/L ²
Minerals	Mg	0.59	mmol/L 0.53-1.06
Electrolytes	Na+	↓ 131.9	mmol/L 138.0-160.0
Electrolytes	K+	↓ 2.8	mmol/L 3.5-5.9
Electrolytes	Na/K	47.6	
Electrolytes	Cl-	104.7	mmol/L 102.7-125.0

Operator:

Comprehensive Diagnosis Panel

QC QC OK

HEM(Hemolysis degree): 0 LIP(Lipemia degree): 0 ICT(Jaundice degree): 2+



Report Explain.

TP



Increase is commonly associated with dehydration and increased globulin. Reduction is commonly associated with blood loss, protein-losing enteropathy, and decreased albumin.

The results only applies to this test sample.

Test Instrument: Mindray vetXpert C5

Time of Printing: 2026-01-15 09:50:46



PETVET CENTRAL VET CLINIC – BARRA
ZONE 4, BARRA, OPOL, MISAMIS ORIENTAL
0927 569 8880

Global Pioneer of Comprehensive Animal Medical Solutions

Better healthcare for all - Since 1991

mindray
animal medical

Biochemistry test report



Patient:	BAILEY	Species:	Canine	Patient ID:	260115001
Client:	SALVA	Gender:	Male	Sample No.:	04
Doctor:		Age:	2Y	Time of analysis:	2026/01/15 09:49



Report Explained.

ALB	↓	Increase is commonly associated with dehydration and corticosteroid administration, etc. Reduction is commonly associated with excessive infusion, malnutrition, hepatic insufficiency or failure, nephropathy, and protein-losing enteropathy.
GLOB	↓	Increase is commonly associated with chronic inflammation and infection, and hyperimmunity, etc. Reduction is commonly associated with insufficient protein intake, anemia, and immunodeficiency.
AST	↑	Increase is commonly associated with liver injury and muscle injury, etc.
ALP	↑	Increase is commonly associated with fracture healing period, hepatobiliary diseases, hyperthyroidism, and osteosarcoma, etc.
GGT	↑	Elevated is commonly associated with bile duct injury or cholestasis, etc.
TBIL	↑	Increase is commonly associated with hemolysis and hepatobiliary dysfunction. Reduction is commonly associated with decreased erythropoiesis, etc.
TBA	↑	Increase is commonly associated with hepatic insufficiency or failure, portal vein shunt, and cholestasis, etc. Reduction is commonly associated with long-term fasting and intestinal malabsorption, etc.
BUN	↑	Increase is commonly associated with high protein diet, gastrointestinal bleeding, nephropathy, and urinary obstruction, etc. Reduction is commonly associated with insufficient protein intake and liver failure, etc.
CK	↑	Increase is commonly associated with trauma, increased muscle activity (such as tetanus and convulsion), myocarditis, and myocardial infarction, etc.
TC	↑	Increase is commonly associated with biliary obstruction, hypothyroidism, hypercorticalismus, nephropathy, diabetes, etc. Reduction is commonly associated with protein loss enteropathy, pancreatic exocrine insufficiency, and hypoadrenocorticism, etc.
TG	↑	Increase is commonly associated with postprandial, obesity, diabetes and hypercorticalismus, etc.
Ca	↓	Increase is commonly associated with hypoadrenocorticism, lymphoma, and nephropathy, etc. Reduction is commonly associated with low calcium diet, hypoalbuminemia, nephropathy, and vitamin D deficiency, etc.
PHOS	↑	Increase is commonly associated with nephropathy, bone healing period, and hyperthyroidism. Decreased in hyperparathyroidism, tumor, etc.
Na+	↓	Increase is commonly associated with salt intoxication, hypertonic NaCl solution rehydration, hyperaldosteronism, and severe dehydration, etc. Reduction is commonly associated with hypoadrenocorticism, diuretic therapy, etc.
K+	↓	Increase is commonly associated with high potassium fluid replacement, diabetes, adrenocortical hypofunction, and acute kidney injury, etc. Reduction is commonly associated with low potassium or potassium-free fluid replacement, vomiting, diarrhea, and hypercorticalismus, etc.

Note: Due to the complexity and individuality of disease diagnosis, the report interpretation is only for your reference. Please consult your doctors for clinical diagnosis results.
The results only applies to this test sample.

Test Instrument: Mindray vetXpert C5

Time of Printing: 2026-01-15 09:50:46



PETVET CENTRAL VET CLINIC – BARRA
ZONE 4, BARRA, OPOL, MISAMIS ORIENTAL
0927 569 8880

Global Pioneer of Comprehensive Animal Medical Solutions

Better healthcare for all - Since 1991

mindray
animal medical